

FUSION 360™ FOLIAR FG-31

1-0-2

Specifically formulated plant constituents designed to contribute to foliar mineral nutrition

GENERAL DESCRIPTION:

For foliar application on various ornamentals, including turf, trees and shrubs, annual and perennial foliage species as well as annual and perennial flowering species.

FUSION 360 FOLIAR FG-31 may also be applied head of stress events, such as increased temperatures above 95°F.

FUSION 360 FOLIAR FG-31 can be applied via boom sprayers, air-blast sprayers or via aerial applications. It can mix with most fungicides, insecticides, adjuvants or nutrients.

CAUTION:

**KEEP OUT OF REACH OF CHILDREN
KEEP OUT OF DIRECT SUNLIGHT & HEAT**

Do Not use **FUSION 360 FOLIAR FG-31** in a dilution less than 8 parts water to 1 part material.

Consult your Fusion 360 technical representative for complete programs, compatible combinations, or rates specific to sensitive plant tissues.

GUARANTEED ANALYSIS:

Total Nitrogen (N)	1.00%
1.0% Nitrate Nitrogen	
Soluble Potash (K ₂ O)	2.00%
Calcium (Ca)	2.20%
Magnesium (Mg)	0.75%
Boron (B)	0.05%
Iron (Fe)	0.10%
Manganese (Mn)	0.06%
Zinc (Zn)	0.12%

Derived from Potassium Nitrate, Boric Acid, Calcium Acetate, Calcium Nitrate, Ferric Nitrate, Magnesium Nitrate, Magnesium Chloride, Manganese Acetate, and Zinc Nitrate

Manufactured By:



2425 Industrial Rowe
Turlock, California 95380
(888) 634-3360
www.fusion360ag.com

**Emergency Telephone Number:
(209) 572-5822**

DIRECTIONS FOR USE:

IF WEATHER IS COOL/OVERCAST OR PLANTS ARE SHOWING SIGNS OF WEAKNESS START WITH REDUCED RATES AND INCREASE OVERTIME.

Stone Fruit & Nut Trees 2-8 gal/100 gal bloom and post-bloom

Citrus & Olives: 1-6 gals/100 gals applied near bloom and post bloom.

Vegetable & Field Crops & Berries: 2-5 gal/100 gal beginning at ~5th leaf stage and continuing throughout the season @ ~7-10 day intervals. **(Note: For berries start with lower rates for initial 2 sprays increasing rates thereafter)**

Transplant Seedlings: 2-5 gal/100 gal beginning at ~2nd-3rd leaf stage and continuing at 4-5 day intervals until field planted.

Vine Crops: 1-6 gal/100 gal pre and post-bloom and continued while active foliage is available. Repeat @ 7-10 day intervals. **(Note: Apply at lower rates for initial 2 sprays increasing rates thereafter)**

Ornamental Trees & Shrubs: 3-8 gal/100 gal applied @ 7-14 day intervals throughout the season.

Alfalfa: 2.5-5 gal/100 gal between cuttings that fall on the 2nd—3rd week of cutting. Minimum of 1 spray between cuttings. 3-4 gal/100 gal of total spray is a useful addition to any spray.

Tropical Fruits: 2-5 gal/100 gal beginning at ~5th leaf stage and continuing throughout the season @ 7-10 day intervals.

LIMITED WARRANTY: Manufacturer or seller makes no warranty, whether expressed or implied, concerning the use of this product other than for the purpose indicated on the label. Neither manufacturer nor seller shall be liable for any injury or damage caused by this product due to misuse, mishandling, or any application not specifically described on this label.

Net Contents:

Lot #

Weight Per Gallon @ 68°F : 11.4 lbs.

_____ 2.5 gallon (9.45 liters)

_____ 250 gallons (946.35 liters)

Information regarding the contents and levels of metals in this product is available on the Internet at www.aapfco.org/metals.html